

## SITE INDEX OF SIAMESE ROSEWOOD (*Dalbergia cochinchinensis* Pierre) IN PLANTATIONS OF THAILAND\*\*

TAENGMOO PHUNCHAISRI<sup>1</sup>, CHONGRAK WACHRINRAT<sup>1\*</sup>, PONTHEP MEUNPONG<sup>1</sup>,  
SUWAN TANGMITCHAROEN<sup>2</sup> AND NAWAPHONG KUASAKUN<sup>1</sup>

<sup>1</sup>Department of Silviculture, Faculty of Forestry, Kasetsart University, Bangkok 10900, Thailand

<sup>2</sup>Forest Research and Development Bureau, Royal Forest Department, Bangkok 10900, Thailand

Received 07 January 2019 / Accepted 16 May 2019

### ABSTRACT

Currently, decreased productivity of Siamese rosewood (*Dalbergia cochinchinensis* Pierre) is a problem due to illegal logging and the guidelines to manage this prevention being may be insufficient. Forest plantations should be promoted to achieve sustainable wood production for this species. Site index (SI) is a measure that can be used to predict forest productivity to identify appropriate sites for planting Siamese rosewood. The objective of the current study was to predict the SI for Siamese rosewood in Thailand. A sample of 78 temporary plots at 26 sites in 16 provinces of Thailand was selected to construct a site index curve with a base age of 30 years. The results showed that the mean age of Siamese rosewood was 28 years (between 13 and 55 years), the mean diameter at breast height (DBH) was 20.03 cm (between 7.99 and 33.80 cm), and the mean dominant height was 20.79 m (between 7.32 and 24.82 m). The SI equation was:  $SI = e(\ln H_{db} - 12.476(A^{-1} - A_b^{-1}))$  with an  $R^2$  value of 0.406 at  $p < 0.0001$  (chosen as 30 years in the present study). The equation indicated that 14, 18, 22, 26 and 30 m trees belonged to site classes I, II, III, IV and V, respectively. Although in natural forest, Siamese rosewood is distributed in dry evergreen and mixed deciduous forests, in the present study, several sites in areas under both *in situ* and *ex situ* conservation were in the good and very good site categories. Thus, both areas can be used to plant Siamese rosewood and increase its productivity.

**Keywords:** Siamese rosewood, site index, temporary plot

### INTRODUCTION

Siamese rosewood (*Dalbergia cochinchinensis* Pierre) is generally found in the dry evergreen and mixed deciduous forests (Veesommai & Kavduengtain 2004) at elevations between 100 and 775 m above sea level (Eiadthong & Tangmitcharoen 2015). This species is native to Thailand, Cambodia, Laos, and Vietnam. Over the years, this species has been severely affected by illegal logging, due to the quality of its hardwood and high durability, leading to a gradual reduction in its numbers (Cadena 2014). It was under the “vulnerable” category and red listed as a threatened species by the International Union for Conservation of Nature

(IUCN) (IUCN 2008), leading to a high demand of its wood and high price (Forest Research and Development Bureau 2010). This has been the prime reason affecting the numbers of Siamese rosewood trees.

According to a survey by the Royal Forest Department of Thailand (2015), the price of Siamese rosewood heartwood was USD 30,000/m<sup>3</sup> on the black market. Such a high price provided sufficient motivation for illegal logging in both natural forest and plantations. In 2014, 1,823 m<sup>3</sup> of Siamese rosewood was seized by the Thai government. The lawsuits filed for illegal logging of Siamese rosewood have increased while productivity has decreased every year in Thailand (Department of National Parks, Wildlife and Plant Conservation 2015). A similar situation exists for other threatened species with similar hardwood characteristics, such as *Dalbergia oliveri* Gamble ex Prain and *Pterocarpus*

\*Corresponding author, e-mail: fforcrw@ku.ac.th

\*\*This paper was presented at the 3<sup>rd</sup> International Conference on Tropical Biology 2018, 20-21 September 2018, Bogor, West Java, Indonesia

*macrocarpus* Kurz (Internal Security Operations Command 2014). The diminishing number of individual trees of such species is of grave concern and needs to be dealt with urgently. Plantations can play an important role in the success of reviving these species. Several economically viable species, including Siamese rosewood, have been promoted in certain areas by the Thai government. Two distinct procedures have been applied: 1) *in situ* conservation areas (conservation of ecosystems and natural habitats and maintenance and recovery of the population that allows the species to exist in the natural environment and in the case of domesticated or cultivated plant species in an environment in which those species have developed special properties); and 2) *ex situ* conservation (conservation of plant genetic resources outside the natural habitat). The current methods being used for selecting areas where this planting Siamese rosewood need to be improved and implemented in the future (Tangmitcharoen 2013).

Site index (SI) is one of the measures used to predict forest productivity, wood volume, and the potential rate of growth (Hanson *et al.* 2002). The relationship between dominant height and age was used to construct a site index curve by Mayhew & Newton (1998). There are several techniques for determining SI such as: (1) the proportion curve or guide curve method; (2) the parameter prediction method; and (3) the different equation method (Clutter *et al.* 1983). In Thailand, the proportion curve method has been used for assessing and predicting the site quality of *Tectona grandis* L.f. (Forestry Research Center 1996) and *Eucalyptus camaldulensis* Dehnh. plantations in the lower northeastern parts

(Insaun *et al.* 2009) and *Acacia mangium* Willd. plantation in Trat province (Jumwong 2006). Siamese rosewood research has also focused on other dimensions such as inventory, ecology, growth and yield. The prediction of SI for Siamese rosewood is yet to be undertaken. Thus, the objective of the current research was to summarize the growth characteristics and develop a SI for Siamese rosewood found in plantations in Thailand.

## MATERIALS AND METHODS

### Study Site and Data Collection

Thailand is a tropical country located between 5°37" N and 20°27" N latitude and 97°22" E and 105°37" E longitude. The annual rainfall ranges between 1,200 and 1,600 mm and the mean annual rainfall is 1,587 mm. The mean annual temperature is around 27° C (Thai Meteorological Department 2014). The study was conducted in Siamese rosewood plantations owned by the Royal Forest Department of Thailand (RFD) and the Forest Industry Organization (FIO). Twenty-six sites were selected from 16 provinces distributed throughout Thailand (Table 1 and Fig. 1). Three temporary sample plots were selected per site (78 temporary sample plots in total). Each temporary plot was 0.16 ha in size and the diameter at breast height over bark (DBH) at 1.30 m above ground and total height (H), of all trees in the plot, were measured. Meteorological data from the Thai Meteorological Department for the sample sites are shown in Table 2.

Table 1 Summary of 26 Siamese Rosewood study sites in Thailand used in the study

No	Plantation	Province	Region	Code	Latitude	Longitude
1	Kam Phaeng Phet Silvicultural Research Station	Kamphaengphet	N	KPP1	16° 33'	99° 30'
2	Kam Phaeng Phet Silvicultural Research Station	Kamphaengphet	N	KPP2	16° 34'	99° 30'
3	Kam Phaeng Phet Silvicultural Research Station	Kamphaengphet	N	KPP3	16° 33'	99° 30'
4	Phit Sa Nu Lok Silvicultural Research Station	Phitsanulok	N	PNL	16° 50'	100° 53'
5	In Tha Khin Silvicultural Research Station	Chiangmai	N	ITK	19° 9'	98° 56'
6	Tha Tum Silvicultural Research Station	Surin	NE	TT	15° 18'	103° 45'
7	Nong Khu Silvicultural Research Station	Surin	NE	NK	14° 41'	103° 45'
8	Huai Tha Silvicultural Research Station	Sisaket	NE	HT1	14° 52'	104° 26'
9	Huai Tha Silvicultural Research Station	Sisaket	NE	HT2	14° 52'	104° 26'
10	Pha Nok Khao Silvicultural Research Station	Khon Kaen	NE	PNK	16° 50'	101° 56'
11	Loei Forestry and Forest Products Research Center	Loei	NE	LOEI	16° 52'	101° 56'

Table 1 (Continued)

12	Udon Tha Ni Forest Nursery Center	Udon Thani Nakhon	NE	UD	16° 50'	102° 56'
13	Mu Si Silvicultural Research Station	Ratchasima Nakhon	NE	MS1	14° 30'	101° 25'
14	Mu Si Silvicultural Research Station	Ratchasima	NE	MS2	14° 30'	101° 25'
15	Ka La Sin Silvicultural Research Station	Kalasin	NE	KLS	16° 28'	103° 24'
16	Dong Lan Silvicultural Research Station	Khon Kaen	NE	DL	16° 49'	101° 59'
17	Huai Rang Plantation	Trat	E	HR	12° 26'	102° 33'
18	Tha Kum Noboru Umeda Plantation	Trat	E	TK1	12° 22'	102° 40'
19	Tha Kum Noboru Umeda Plantation	Trat	E	TK2	12° 23'	102° 39'
20	Tha Kum Noboru Umeda Plantation	Trat	E	TK3	12° 23'	102° 39'
21	Rat Cha Bu Ri Silvicultural Research Station	Ratchaburi	C	RBR	13° 34'	99° 44'
22	Central Silvicultural Research center	Kanchanaburi	C	PK1	14° 24'	98° 55'
23	Central Silvicultural Research center	Kanchanaburi	C	PK2	14° 24'	98° 55'
24	Thong Pha Phum Silvicultural Research Station	Kanchanaburi	C	TPP	14° 42'	98° 39'
25	Southern Silvicultural Research center	Songkhla Prachuap Khiri Khan	S	PT	7° 1'	100° 17'
26	Sai Thong Silvicultural Research Station	Khan	S	ST	10° 59'	99° 27'

Note: Regions as identified by the Thai Meteorological Department are: “N” is north, “NE” is northeast, “E” is east, “C” is central, and “S” is south.

Table 2 Meteorological data for the study sites

No.	Station name	Average temperature (°C)	Average relative humidity (%)	Rainfall (mm/year)	Study site
1	Phitsanulok	28.1	74.2	1,399	PNL
2	Kamphaengphet	28.5	76.8	1,372	KPP1, KPP2, KPP3
3	Chiangmai	27.2	71.3	1,158	ITK
4	Pakchong Agromet	26.4	73.3	1,206	MS1, MS2
5	Thatom	27.9	74.1	1,313	TT
6	Surin	27.7	72.4	1,454	NK
7	Sisaket Agromet	27.7	74.8	1,420	HT1, HT2
8	Loei	26.9	73.2	1,310	LOEI
9	Kalasin	27.2	72.0	1,340	KLS
10	Khon Kaen	27.7	70.4	1,196	PNK, DL
11	Udonthani	27.5	71.1	1,338	UD
12	Trat	27.9	81.4	4,988	HR, TK1, TK2, TK3
13	Ratchaburi	28.6	78.1	1,067	RBR
14	Kanchanaburi	29.0	70.4	1,104	PK1, PK2
15	Thong Phaphum	28.1	78.5	1,827	TPP
16	Prachuap Khiri Khan	28.8	76.3	1,056	ST
17	Hatyai	28.3	79.4	1,793	PT

**Determination of SI Equation**

The height of the tree is important in the creation of the SI but the mean height of a stand is usually sensitive to age, site, class and stand density. Thus, the dominant height is normally used to represent a stand and specifically mean height of the 100 tallest trees/ha is most widely used to construct an SI (Alder 1980). In this case, the SI equation was constructed using the top 16 tallest trees per plot (plot size of 0.16 ha). There are several techniques for determining the

SI. However, the current study only had access to data derived from temporary plots and consequently, the guide curve method was used (Alder1980) based on equation (1) and the SI equation was constructed using equation (2).

$$\ln (H_{do}) = \ln b_0 + b_1 A^{-k} \dots \dots \dots (1),$$

$$SI = e (\ln (H_{do}) + b_1 (A^{-1} - A_b^{-1})) \dots \dots \dots (2),$$

where  $H_{do}$  is the dominant height,  $\ln$  is the natural logarithm,  $b_0$  and  $b_1$  are regression coefficients,  $A$  is the stand age,  $k = 1$ ,  $SI$  is the site index,  $e$  is the natural exponent, and  $A_b$  is

the base age (chosen as 30 years in the present study).

Based on equation 2, 5 site classes were used (I, II, III, IV, and V) indicating very poor, poor, moderate, good, and very good sites, respectively. The upper and lower bounds for each SI class were determined using simple interpolation and extrapolation in the Excel software package (Microsoft Corp.; Redmond, CA, USA).

## RESULTS AND DISCUSSION

The growth characteristics of Siamese rosewood, as determined from 26 sites, are presented in Table 3. The mean age was 28 years (between 13 and 55 years), mean DBH was 20.03 cm (between 7.99 and 33.80 cm), mean H

was 16.48 m (ranging between 6.18 and 23.20 m), and the mean  $H_{do}$  was 20.79 m (ranging between 7.32 and 24.82 m). The mean DBH, H and  $H_{do}$  of the TT plantation were lowest (see Table 1 for identification of the plantation abbreviations used). Both the highest mean DBH and mean H were in the KLS plantation and the highest mean of  $H_{do}$  was in the PNL plantation. The mean dominant heights for KLS and PNL were similar (24.18 and 24.82 m, respectively) despite their stand ages being rather different (42 and 27 years, respectively). This could be interpreted as the younger stand age alone indicating site suitability which is not necessarily correct. Thus, the methodology for assessing the appropriate site using the site index had to decrease the bias associated with the different-aged stands.




Figure 1 Location of Siamese rosewood study sites in Thailand

Table 3 Growth characteristics of the Siamese rosewood in plantations located at the 26 sites in Thailand

No	Site code	Age (years)	DBH ± SD (cm)	H ± SD (m)	$H_{do}$ ± SD (m)
1	TT	13	7.99 ± 3.94	6.18 ± 2.65	7.32 ± 2.22
2	RBR	22	23.42 ± 7.48	14.06 ± 3.30	14.55 ± 3.00
3	NK	25	13.23 ± 5.66	10.78 ± 3.85	16.03 ± 2.40
4	HT1	25	18.38 ± 6.06	14.58 ± 3.56	17.55 ± 1.49
5	HT2	25	21.02 ± 8.27	16.64 ± 4.13	17.43 ± 3.03
6	PNK	54	31.59 ± 7.16	22.64 ± 4.83	22.92 ± 4.70
7	LOEI	55	18.34 ± 6.07	17.42 ± 4.68	22.14 ± 2.43
8	UD	20	14.18 ± 5.73	13.82 ± 4.95	17.50 ± 4.13

Table 3 (Continued)

9	KPP1	23	18.43 ± 7.30	16.72 ± 3.93	20.62 ± 1.90
10	MS1	28	21.15 ± 6.15	16.74 ± 3.18	20.02 ± 1.16
11	MS2	28	15.41 ± 6.29	13.96 ± 4.63	20.72 ± 2.38
12	HR	30	20.49 ± 7.92	16.21 ± 5.57	22.72 ± 2.61
13	TK1	32	25.06 ± 10.62	17.58 ± 6.16	23.53 ± 2.52
14	TK2	35	22.21 ± 8.48	15.82 ± 5.36	22.04 ± 1.74
15	TK3	36	25.40 ± 9.18	15.09 ± 5.25	21.38 ± 2.01
16	KLS	42	33.80 ± 9.92	23.20 ± 5.02	24.18 ± 4.94
17	PK2	18	18.39 ± 7.17	15.69 ± 4.64	20.71 ± 1.74
18	PK1	21	16.99 ± 7.65	15.09 ± 4.98	20.72 ± 1.53
19	KPP2	23	21.11 ± 7.07	19.47 ± 5.13	23.86 ± 2.38
20	KPP3	23	20.65 ± 9.55	17.54 ± 6.68	24.35 ± 1.96
21	PT	26	20.29 ± 8.14	18.24 ± 5.45	23.08 ± 0.90
22	PNL	27	15.97 ± 6.83	17.93 ± 5.45	24.82 ± 1.22
23	TPP	27	25.55 ± 11.14	20.01 ± 5.26	24.30 ± 1.28
24	ST	28	19.18 ± 7.67	19.64 ± 4.62	24.42 ± 1.15
25	DL	16	16.66 ± 5.68	16.06 ± 3.42	19.89 ± 1.92
26	ITK	20	15.85 ± 6.89	17.30 ± 5.16	23.60 ± 2.16

The SI equation for Siamese rosewood obtained using the proportional curve method at a base age of 30 years was  $SI = e(\ln H_{do} - 12.476(A^{-1} - A_b^{-1}))$  with an  $R^2$  value of 0.406 and  $p < 0.0001$ . Five stratification levels were used based on site classes I through V for H values of 14, 18, 22, 26 and 30 m, respectively. Figure 2 shows that a 30 years Siamese rosewood tree with a dominant height of 22 m belonged to site class III. This stand was predicted as a moderate site for planting. Likewise, we also determined the best site planting (class V) for which trees had to be 30 m tall at age 30 years. In contrast, trees on poorer sites (site I) were only 14 m tall at age 30 years.

The number of study sites in the classifications were 1, 6, 9, 8, and 2 for SI classes I, II, III, IV, and V, respectively (Table 3). The lower and upper bounds of a very poor site class were 12.00 and 15.99 m, respectively, thus, the TT plantation was in this category. On the other hand, the lower and upper bounds of a very good site were 28.00 and 32.00 m, respectively, and the DL and ITK plantations were in this

category. In this study, most of the measured sites were in either the moderate or good site classes. Even though the growth rate based on DBH, H, and  $H_{do}$  of KLS and PNL were satisfactory, the results showed that DL and ITK were very good sites for Siamese rosewood compared at the base age of 30 years.

Kliangpibool (2012) reported the suitable age for Siamese rosewood logs was between 18 and 25 years at the price was USD 530 to 1,000/ha. However, the valuation of Siamese rosewood does not depend only on the quantity as the economic valuation of hardwood is important too. Kuasakun (2017) determined that the heartwood percentage of Siamese rosewood at the Mu Si Silvicultural Research Station (MS) was 68.30% while the SI for MS1 and MS2 were moderate site. In contrast, the heartwood percentage of Siamese rosewood at In Tha Kin Silvicultural Research Station (ITK) was 21.97 while the SI of ITK in the present study indicated it was a very good site; nevertheless, the increasing heartwood should result with increasing age as well (Visaratana *et al.* 2016).




Figure 2 SI curves for Siamese rosewood in Thailand

Note: Dotted lines indicate the upper and lower bounds for each class based on simple interpolation/extrapolation.

Table 3 Site class, SI and bounds at base age 30 years of Siamese rosewood in Thailand, where the last column indicates the class according to site quality

Site class	SI	Number of study sites	Lower limit (m)	Upper limit (m)	Site code
I (very poor)	14	1	12.00	15.99	TT
II (poor)	18	6	16.00	19.99	RBR, NK, HT1, HT2, PNK, LOEI
III (moderate)	22	9	20.00	23.99	UD, KPP1, MS1, MS2, HR, TK1, TK2, TK3, KLS
IV (good)	26	8	24.00	27.99	PK2, PK1, KPP2, KPP3, PT, PNL, TPP, ST
V (very good)	30	2	28.00	32.00	DL, ITK




Figure 3 *In situ* study sites of Siamese rosewood in northeastern and eastern Thailand

The natural distribution of Siamese rosewood is in the dry evergreen and mixed deciduous forests where located northeastern and eastern Thailand (Fig. 3). Both regions have *in situ* conservation areas in which were located the study sites HR, TK1, TK2 and TK3 in the eastern, and TT, NK, HT1, HT2, PNK, LOEI, UD, MS1, MS2, KLS and DL in the northeastern regions. Several sites in the *in-situ* conservation areas were not as appropriate as plantation sites as in the *ex-situ* conservation areas. The better *ex situ* sites were PK1, PK2, KPP1, KPP2, PT, PNL, TPP, ST and ITK were determined to be in the good and very good site classes (IV and V, see Fig. 4). Suanpaga & Boonyuan (2016) compared the growth of eight appropriate tree species for planting in the Songkhla province, southern Thailand (an *ex-situ* conservation area for Siamese rosewood). Based on their results, they recommended Siamese rosewood and three other species (*Dipterocarpus alatus* Roxb, *Xylia xylocarpa* (Roxb.) Taub. var. *kerrii* (Craib & Hutch.) I. C. Nielsen and *Azadirachta excelsa* (Jack) Jacobs).

Illegal logging has continued (Sudsakorn 2012) and so plans to increase sustainable productivity need to consider prevention and promotion. Chaipunya (2016) considered three aspects to maintain sustainable production from current stands: preventing illegal rosewood logging, preventing transportation, and preventing export of logs or wood products. In addition, Visaratana *et al.* (2016) presented a case to promote both forest management and silvicultural techniques via plantations located on good sites to provide increasing productivity and economic value. Siamese rosewood can fix nitrogen through *Rhizobium* bacteria that can increase the health of ecological systems by fixing nutrients for use also by other species (Mungklarat 2001) and to improve soil fertility for reforestation purposes (So *et al.* 2010). With regard to the current study, several sites in the *ex-situ* conservation area belonged to either the good or very good site classes. Thus, both *ex-situ* and *in-situ* conservation areas can be used for plantation purposes to increase productivity.




Figure 4 *Ex situ* study sites of Siamese rosewood in northern, central, and southern Thailand

## CONCLUSION

A site index curve and equation are necessities for Thailand forestry to support Siamese rosewood plantations dating back more than 50 years; however, extensive research commenced only recently. As a result, knowledge is still lacking of the site requirements for plantations of this species. Based on the growth characteristic of Siamese rosewood from the Royal Forest Department of Thailand (RFD) and the Forest Industry Organization (FIO) plantations at 26 sites represented the age ranges between 13 and 55 years, the DBH ranges between 7.99 and 33.80 cm, the H ranges between 6.18 and 23.20 m and the  $H_{do}$  ranges between 7.32 and 24.82 m. In addition, five SI classes were constructed with most sites in the moderate site class. The SI equation developed in this study can be applied to help avoid planting of Siamese rosewood on inappropriate lower quality sites and place more emphasis on identifying higher quality sites for plantation establishment. Several sites in the *ex-situ* conservation area belonged either to the good or very good site classes. Thus, both *ex-situ* and *in-situ* conservation areas can be used for plantation purposes to increase productivity.

## ACKNOWLEDGEMENTS

This research was supported in part by a Graduate Program Scholarship from the Graduate School, Kasetsart University, Bangkok, Thailand. We thank the Royal Forest Department of Thailand (RFD) and the Forest Industry Organization (FIO) for helping with the supply of data and assistance with fieldwork.

## REFERENCES

- Alder D. 1980. Forest volume estimation and yield prediction vol. 2 yield production. Italy: Food and Agriculture Organization of United Nation.
- Cadena A [Internet]. 2014. (Un)protected Areas: Rosewood poaching in Thailand's Dong Phrayayen-KhaoYai world heritage site: IUCN.com; [updated 2014 Feb 24; cited 2018 Jun 20]. Available from: <https://www.iucn.org/content/unprotected-areas-rosewood-poaching-thailand%E2%80%99s-dong-phrayayen-khao-yai-world-heritage-site>

- Chaipunya M. 2016. Monitoring of ad hoc plan's implementation on combating illegal logging of rosewood in Northeastern protected area 2013. Bangkok (TH): Department of National Parks, Wildlife and Plant Conservation.
- Clutter JL, Fortson JC, Pienaar LV, Brister GH, Bailey RL. 1983. Timber Management, a Quantitative Approach. New York (US): John Wiley & Sons, Inc.
- Department of National Parks, Wildlife and Plant Conservation. 2015. Special case to solve the problem in the area of Thap Lan National Park and Pang Sida National Park. Bangkok (TH): Department of National Parks, Wildlife and Plant Conservation.
- Eiadthong W, Tangmitcharoen S. 2015. Ecological and geographical distributions and its demographic status of Siamese rosewood (*Dalbergia cochinchinensis* Pierre) in Thailand. In: FORCONS proceeding 2015.
- Forestry Research Center. 1996. Exploration and yield assessment of the Forest Industry Organization. Bangkok (TH): Faculty of Forestry, Kasetsart University.
- Forest Research and Development Bureau. 2010. General characteristics of Thai timbers. Bangkok (TH): The Royal Forest Department of Thailand.
- Hanson EJ, Azuma DL, Hiserote BA. 2002. Site index equations and mean annual increment equations for Pacific Northwest Research Station forest inventory and analysis inventories, 1985-2001. Res. Note PNW-RN-533. Portland (OR, US): Department of Agriculture, Forest Service, Pacific Northwest Research Station. p. 1-24.
- Insaun W, Wachrinrat C, Srigongpan R. 2009. Site quality assessment for *Eucalyptus camaldulensis* Dehnh. plantation in the lower part of Northeastern Thailand. Thai J For 28:13-27.
- Internal Security Operations Command. 2014. Model scheme of forest protection, land invasion of the state and sustainable management of natural resources. Bangkok (TH): Internal Security Operations Command.
- International Union for Conservation of Nature [Internet]. 2008. IUCN Red list of threatened species. Version 2008. [cited 2018 Jun 20]. Available from: <http://www.iucnredlist.org/details/32625/0>
- Jumwong N. 2006. Site potential evaluation of *Acacia mangium* Willd. plantation area in Trat province, Thailand [Dissertation]. Retrieved from Kasetsart University.
- Kliangpibool N. 2012. *Dalbergia cochinchinensis* Pierre plantation. Bangkok (TH): The Royal Forest Department of Thailand.

- Kuasakun N. 2017. Growth and heartwood ratio of Siamese rosewood (*Dalbergia cochinchinensis* Pierre) in various plantations [Dissertation]. Retrieved from Kasetsart University.
- Mayhew JE, Newton AC. 1998. The silviculture of mahogany. England: CABI Publishing.
- Mungklarat J. 2001. The establishment of *Dalbergia cochinchinensis* Pierre plantation. Bangkok (TH): Royal Forest Department of Thailand.
- Royal Forest Department of Thailand. 2015. RFD news. Bangkok (TH): Royal Forest Department of Thailand.
- So T, Theilade I, Dell B. 2010. Conservation and utilization of threatened hardwood species through reforestation - An example of *Azelia xylocarpa* (Kruz.) Craib and *Dalbergia cochinchinensis* Pierre in Cambodia. *Pacific Conserv Biol* 16:101-16.
- Suanpaga W, Boonyuan S. 2016. Growth and aboveground biomass of eight tree species in species trial for appropriate planting in Had Yai District, Songkhla Province. In: FORCONS proceeding 2016. p. 136.
- Sudsakorn C. 2012. Educational guidelines for preventing and combating an illegal logging of Pa Yung (*Dalbergia cochinchinensis*) [Dissertation]. Retrieved from Joint War College.
- Tangmitcharoen S. 2013. Siamese rosewood: Conservation and management of genetic resources. In: Royal Forest Department Proceedings 2013. Forestry meeting on production and forest research towards sustainable development.
- Thai Meteorological Department [Internet]. 2014. The climate of Thailand; [cited 2018 Sep 24]. Available from: [https://www.tmd.go.th/info/climate\\_of\\_thailand-2524-2553.pdf](https://www.tmd.go.th/info/climate_of_thailand-2524-2553.pdf)
- Veesommai U, Kavduengtain P. 2004. Wild trees in Thailand 1. Bangkok (TH): H.N. Group Co., Ltd.
- Visaratana T, Pinthong A, Hutasangchai C. 2016. Estimation heartwood of *Dalbergia cochinchinensis* Pierre in the plantation. In: FORCONS proceeding 2016. p. 72.