

THE RANGE EXPANSION OF *Parachromis managuensis* GÜNTHER, 1867 (PERCIFORMES, CICHLIDAE) IN JAVA, INDONESIA

MAHENO SRI WIDODO¹, VERYL HASAN^{2*}, ABDUL RAHEM FAQIH¹, MAFTUCH¹,
R. ADHARAYAN ISLAMY¹ AND FELIPE POLIVANOV OTTONI³

¹Department of Aquatic Resources Management, Universitas Brawijaya, Malang 65145, Indonesia

²Department of Aquaculture, Universitas Airlangga, Surabaya 60115, Indonesia

³Laboratório de Sistemática e Ecologia de Organismos Aquáticos, Centro de Ciências Agrárias e Ambientais, Campus Universitário, Universidade Federal do Maranhão, BR-222, KM 04, S/N, Boa Vista, CEP 65500-000, Brazil

Received 28 June 2019 / Accepted 24 May 2021

ABSTRACT

The Jaguar cichlid, *Parachromis managuensis* (Günther, 1867), is native to Central America, with introductions reported from West Java and Central Java Provinces of Indonesia. On 7-8 January 2019, sixteen specimens of *P. managuensis* were collected from Karangates, the largest hydropower reservoir in East Java Province of Indonesia. A description of the morphological characters of specimens is provided.

Keywords: Cichlid, distribution, freshwater fish, Jaguar Guapote

INTRODUCTION

Parachromis managuensis (Günther, 1867), is a cichlid native to Costa Rica, Nicaragua and Honduras (Conkel 1993), but it has been introduced to several countries in North America (Fuller *et al.* 1999), South America (Magalhães & Jacobi 2013), and Southeast Asia (Agasen *et al.* 2006). *Parachromis managuensis* exhibits highly predatory habits and tolerance to new habitats (Rosana *et al.* 2006; Agasen *et al.* 2006), which make *P. managuensis* potential to become an invasive species (Yamamoto & Annete 2000).

Parachromis managuensis is generally sold as ornamental fish and has not been cultured openly. *P. managuensis* in Indonesia is firstly found in the freshwaters of West Java (Dahrudin *et al.* 2016) and Central Java (Hedianto *et al.* 2013) Provinces of Indonesia.

The presence of *P. managuensis* at the Karangates Reservoir is considered a new finding because there has been no previous record of exotic fish culture in Karangates Reservoir, the largest hydropower reservoir in East Java Province.

MATERIALS AND METHODS

The Fish Sampling and Description of the Study Sites

Sixteen (16) live specimens of *P. managuensis* were obtained from a local angler during fieldwork conducted on 7 - 8 January 2019 at the Karangates Reservoir (8°11'16"S; 112°27'22"E) (Fig. 1). Administratively, the Karangates Reservoir is located in Malang Regency, East Java Province, Indonesia. Fishing gear used by the angler was a medium hook with a bottom and using worms as bait (Stein *et al.* 2012).

*Corresponding author, email: veryl.hasan@fpk.unair.ac.id



Figure 1 The collecting point of *Parabromis managuensis* at the Karangates Reservoir in East Java Province

Fish Identification

The morphological characters of the fish specimens were determined by using the methods employed by Kullander and Hartel (1997) and Bussing (1998).

RESULTS AND DISCUSSION

Specimens Collection

The sixteen (16) live specimens of *P. managuensis* had a range of total length between 9.9 mm and 26.6 cm. Five (5) of the specimens were preserved in 96% alcohol solution (Hasan & Taman 2019) and transported to the Hydrobiology Laboratory, Universitas Brawijaya, Malang, Indonesia (voucher no. Hb.Pm.I.2019). The remaining eleven (11) specimens were kept as livestock at the Fish Reproduction Laboratory, Universitas Brawijaya, Malang, Indonesia. The 11 living specimens were transported in oxygen-filled polyethylene bags.

Diagnosis

The morphological characters of the specimens are as follows: a large mouth, projecting lower jaw, prominent enlarged canine teeth, a more or less continuous black stripe between the eye and opercular margin, and another stripe between the eye and the lower angle of the opercle, and a row of black blotches along the middle of the side.

The fish can be distinguished from other members of the genus by having the expanded preopercle at the angle. It has silvery or golden-green to purple body colors and black spots on the fins and body. There are also numerous black spots on the anal and caudal fins. The fish has moss green back, purple iridescence sides, and a whitish or yellowish belly. It also has whitish yellowish, or blue iridescence dorsal interspaces, and a black blotch on the caudal-fin base.

All of these characteristics were found in every specimen collected from the Karangates Reservoir, East Java Province, Indonesia (Fig. 2).



Figure 2 Specimen of *Parachromis managuensis* captured on 8 January 2019 in Karangkates Reservoir, East Java Province

Distribution

The discovery of *P. managuensis* in Karangkates Reservoir is the first record of this species in East Java Province. Other discoveries occurred at reservoirs in West Java and Central Java Provinces. The discovery in Karangkates

Reservoir represents a discovery in the eastern part of Java Island, which is around 400 km apart from the Central Java Province (Fig. 3). This record is an important contribution to understanding the dispersal of alien fish species in Indonesia.




Figure 3 A. Distribution of *Parachromis managuensis* in Java Island

Notes: Red square: West Java Province; Blue square: Central Java Province; Green square: East Java Province).

B. Location of the Karangkates Reservoir in East Java Province

Note: The green square indicates the new record of *P. managuensis*

We speculated that *P. managuensis* were released into Karangates Reservoir in East Java Province by exotic fish hobbyists without clear purposes. Further investigation is warranted to determine the source of *P. managuensis* in East Java Province because the reservoir has never been used for any exotic fish culture industry. Control and prevention of further introductions are needed to prevent alien fish from disturbing the freshwaters ecosystem (Hasan *et al.* 2020; Wijayanti *et al.* 2021; Hasan *et al.* 2021).

CONCLUSION

Parachromis managuensis is a non-native fish of Indonesia. This fish species has been found in freshwater in West Java and Central Java Provinces and also in the Karangates Reservoir in East Java Province. The existence of *P. managuensis* in East Java is considered a new finding and added data on the alien fish species distribution in Indonesia.

ACKNOWLEDGMENTS

The authors sincerely thank the Ministry of Finance of Indonesia for funding this study (Grant no. 20160221035555) and the local angler as our field guide.

REFERENCES

- Agasen EV, Clemente JP, Rosana MR, Kawit NS. 2006. Biological investigation of Jaguar Guapote *Parachromis managuensis* (Günther, 1867) in Taal Lake, Philippines. *J Environ Sci Manag* 9(2):20-30.
- Barros LC, Santos U, Zanuncio JC, Dergam JA. 2012. *Plagioscion squamosissimus* (Sciaenidae) and *Parachromis managuensis* (Cichlidae): A threat to native fishes of the Doce River in Minas Gerais, Brazil. *PLoS ONE* 7(6): e39138. <https://doi.org/10.1371/journal.pone.0039138>.
- Bussing WA. 1998. *Peces de Las Aguas Continentales de Costa Rica (Freshwater Fishes of Costa Rica)*. 2nd edition. Editorial de la Universidad de Costa Rica, San José.
- Conkel D. 1993. *Cichlids of North and Central Americas*. New York (US): TFH Publications. 191 p.
- Dahrudin H, Hutama A, Busson F, Sauri S, Hanner R, Keith P, ..., Hubert N. 2016. Revisiting the ichthyofauna diversity of Java and Bali through DNA barcodes: Taxonomic coverage, identification accuracy, cryptic diversity and identification of exotic species. *Mol Ecol Resour* 17(2):288-99. <https://doi.org/10.1111/1755-0998.12528>.
- Hasan V, Widodo, MS, Islamy RA, Pebriani DAA. 2020. New records of alligator gar, *Atractosteus spatula* (Actinopterygii: Lepisosteiformes: Lepisosteidae) from Bali and Java, Indonesia. *Acta Ichthyologica et Piscatoria* 50(2): 233-236. <https://doi.org/10.3750/AIEP/02954>.
- Hasan V, Tamam MB. 2019. First record of the invasive Nile Tilapia, *Oreochromis niloticus* (Linnaeus, 1758) (Perciformes, Cichlidae), on Bawean Island, Indonesia. *Check List* 15 (1): 225-227. <https://doi.org/10.15560/15.1.225>.
- Wijayanti A, Hasan V, Tamam MB. 2021. Range expansion of *Oreochromis niloticus* (Linnaeus, 1758) (Perciformes, Cichlidae) in Java Sea and first record for Masalemba Island. *IOP Conference Series Earth and Environmental Science* 718(1): 012096.
- Hasan V, Valen FS, Islamy RA, Widodo MS, Saptadajaja AM, Islam I. 2021. Short Communication: Presence of the vulnerable freshwater goby *Sicyoptus auxilimentus* (Gobiidae, Sicydiinae) on Sangehe Island, Indonesia. *Biodiversitas* 22: 571-9 <https://doi.org/10.13057/biodiv/d220208>.
- Hedianto DA, Purnomo K, Warsa A. 2013. Interactions of food resources utilization by fish communities in Penjalin Reservoir, Central Java. *Bawal* 5(1): 33-40.
- Fuller PL, Nico LG, Willians JD. 1999. Nonindigenous fish introduced into inland waters of the United States. Bethesda (US): American Fisheries Society.
- Gestring KB, Shafland PL. 1997. Status and selected life history attributes of the exotic Jaguar Guapote (*Cichlasoma managuense*) in Florida. *Florida Scientist* 60(3):137-42.
- Kwik JTB, Kho ZY, Quek BS, Tan HH, Teo DCJ. 2013. Urban stormwater ponds in Singapore: Potential pathways for spread of alien freshwater fishes. *BioInvasions Rec* 2(3):239-45. <http://dx.doi.org/10.3391/bir.2013.2.3.11>.
- Kullander SO, Hartel KE. 1997. The systematic status of cichlid genera described by Louis Agassiz in 1859: *Amphiphobus*, *Baiodon*, *Hypsophrys* and *Parachromis* (Teleostei: Cichlidae). *Ichthyol Explor Freshw* 7:193-202.
- Magalhães ALB, Jacobi CM. 2013. Invasion risks posed by ornamental freshwater fish trade to southeastern Brazilian rivers. *Neotrop Ichthyol* 1(3):433-41. <https://doi.org/10.1590/S1679-62252013005000003>.

- Page LM, Burr BM. 1991. A field guide to freshwater fishes of North America and North of Mexico. Boston (US): Houghton Mifflin Company. 432 p.
- Rosana MR, Agasen EV, Villanueva LS, Clemente Jr JP, Kawit NS, de la Vega JT. 2006. Status and economic impact of *Parachromis Maraguensis* in Taal Lake, Philippines. *J Environ Sci Manag* 9(2):1-19.
- Sampaio WMS, Belei F, Giongo P, Dergam JA, Orsi ML. 2017. *Heterotilapia buttikoferi* (Hubrecht, 1881) (Perciformes: Cichlidae), an introduced exotic fish in the upper Paraná river basin. *Check List* 13(4): 245-250. <https://doi.org/10.15560/13.4.245>.
- Stein JA, Shultz AD, Cooke SJ, Danylchuk AJ, Hayward K, Suski CD. 2012. The influence of hook size, type, and location on hook retention and survival of angled bonefish (*Albula vulpes*). *Fish Res* 113:147-52.
- Yamamoto MN, Annete WT. 2000. Hawaii's Native and Exotic Freshwater Animals. Honolulu (US): Mutual Publishing.